

First-in-human study of RLY-2608, a pan-mutant and isoform selective PI3K α inhibitor, as a single agent in patients with advanced solid tumors and in combination with fulvestrant in patients with advanced breast cancer

Cristina Saura,¹ Lucía Sanz Gómez,¹ Janice S. Kim,² Komal Jhaveri,³ Angel I. Guerrero Zotano,⁴ Pablo Tolosa,⁵ Anne Schott,⁶ Giuseppe Curigliano,⁷ Cesar A. Perez,⁸ Rita Nanda,⁹ Erika P. Hamilton,¹⁰ Jason T. Henry,¹¹ Kari B. Wisinski,¹² Sarah L. Sammons,¹³ Jennifer Segar,¹⁴ Santiago Ponce-Aix,¹⁵ Milana Bergamino,¹⁶ Alexander I. Spira,¹⁷ Alison M. Schram,³ Andreas Varkaris²

¹Vall d'Hebron University Hospital, Vall d'Hebron Institute of Oncology, Barcelona, Spain; ²Massachusetts General Hospital, Boston, Massachusetts, USA; ³Memorial Sloan Kettering Cancer Center, New York, New York, USA; ⁴Instituto Valenciano de Oncología, Valencia, Spain; ⁵Hospital Universitario 12 de Octubre, Madrid, Spain; ⁶The University of Michigan, Ann Arbor, Michigan, USA; ⁷Istituto Europeo di Oncologia, IRCCS, University of Milano, Milan, Italy; ⁸Sarah Cannon Research Institute /Florida Cancer Specialists, Orlando, Florida, USA; ⁹The University of Chicago Medical Center (UCCM), Chicago, Illinois, USA; ¹⁰Sarah Cannon Research Institute, Nashville, Tennessee, USA; ¹¹Sarah Cannon Research Institute/HealthOne Denver, Denver, Colorado, USA; ¹²University of Wisconsin, Madison, Carbone Cancer Center, Madison, Wisconsin, USA; ¹³Dana-Farber Cancer Institute, Boston, Massachusetts, USA; ¹⁴University of Arizona Cancer Center, Tucson, Arizona, USA; ¹⁵Institut Gustave Roussy, Villejuif, France; ¹⁶Institut Catalan d'Oncologia Badalona (ICO), Barcelona, Spain; ¹⁷Virginia Cancer Specialists, Fairfax, Virginia, USA


BACKGROUND

- Oncogenic mutations in *PIK3CA* are common driver events in solid tumors, particularly in breast cancer (BC), where mutations occur in up to ~35% of HR+, HER2– advanced BC patients^{1,2}
- In BC, *PIK3CA* mutations in the kinase (i.e. H1047R) and helical (i.e. E542K, E545K) domains are the most common and contribute to resistance to endocrine therapy
- The approval of orthosteric PI3K α inhibitors, like alpelisib (approved for HR+, HER2– advanced BC), validates PI3K α as a therapeutic target. However, these inhibitors also potentially inhibit WT PI3K α , leading to significant hyperglycemia and rash, which limit dose intensity and potential for use in combination regimens³
- A PI3K α mutant-selective inhibitor could avoid WT PI3K α toxicity and potentially improve dose intensity, efficacy and tolerability, and be used in combinations, benefiting broader populations of patients

RATIONALE


- RLY-2608 is the first oral, pan-mutant- and isoform-selective allosteric inhibitor of PI3K α (Figure 1a)
- RLY-2608 demonstrates biochemical selectivity for both kinase and helical mutants over WT PI3K α (Figure 1b) and kinome selectivity, including over other PI3K isoforms

Figure 1. Novel allosteric mechanism of RLY-2608 enables selectivity for PI3K α mutants⁴



- BC xenograft models with *PIK3CA* kinase and helical mutations demonstrate anti-tumor activity of RLY-2608 monotherapy, while maintaining glucose homeostasis (Figure 2)
- We initiated the first-in-human ReDiscover study (NCT05216432) to evaluate the safety and clinical activity of RLY-2608 as a single agent in patients with advanced solid tumors with *PIK3CA* mutations and in combination with fulvestrant and fulvestrant + a CDK4/6 inhibitor in patients with *PIK3CA*-mutant, HR+, HER2– metastatic BC
- To date, RLY-2608 plus fulvestrant has completed dose escalation; RP2Ds have been identified based on safety, PK, PD, and anti-tumor activity, and are being investigated in the combination regimens

Figure 2. RLY-2608 leads to tumor growth inhibition in HR+ BC mouse xenograft models with *PIK3CA* kinase and helical mutations while maintaining glucose homeostasis⁵




Dotted horizontal line: starting tumor size
*This model also carries a second mutation at K567R; †HSC2 model; ‡Juric D, et al. *J Clin Oncol.* 2018;36:1291–1299 and Juric D, et al. Presented at San Antonio Breast Cancer Symposium 2019, San Antonio, TX, USA. Abstract OT1-08-04; poster P5-17-05.


STUDY DESIGN

This is a global, multi-center, dose-escalation/-expansion study of RLY-2608 as a single agent in adults who have advanced solid tumors (refractory or intolerant to, or who decline, standard therapy) and RLY-2608 in combination with fulvestrant ± ribociclib in previously treated patients with HR+, HER2– metastatic BC (Figure 3)

Figure 3. Study design



RLY-2608 PO BID or QD



- RLY-2608 is administered PO QD in 4-week cycles
- AEs per CTCAE v5, PK, biomarkers (pAKT, mutant ctDNA and insulin pathway markers) and anti-tumor activity are assessed serially
- Dose escalation employs a BOIN design to identify the MTD. Following dose escalation, expansion cohorts are enrolling patients with select *PIK3CA*-mutated solid tumors for treatment with RLY-2608 monotherapy and patients with HR+, HER2– metastatic BC for treatment with RLY-2608 plus fulvestrant ± ribociclib

*Includes *PIK3CA*mut clear cell ovarian cancer, *PIK3CA*mut HNSCC, and *PIK3CA*mut cervical cancer; *Triple combination arms will include the CDK4/6 inhibitor ribociclib.

Eligibility criteria

- ≥18 years of age
- ≥1 documented primary oncogenic *PIK3CA* mutation per local assessment (tumor or blood)
- Eastern Cooperative Oncology Group performance status 0–1
- Evaluable disease per RECIST v1.1
- No prior PI3K α inhibitor*
- For the RLY-2608 combination and triplet arms, patients must have previous treatment with ≤1 chemotherapy, ≥1 CDK4/6 inhibitor, and ≥1 anti-estrogen therapy, and no prior treatment with PI3K α , AKT, or mTOR inhibitors*
- No Type 1 or Type 2 diabetes requiring antihyperglycemic medication, or fasting plasma glucose ≥140 mg/dL and HbA1c ≥7.0%

*Except Part 2 RLY-2608 + fulvestrant combination group intolerant to PI3K α inhibitors.

ENROLLMENT

- The target enrollment for RLY-2608 is 400 patients. Recruitment is ongoing in 24 study centers in the USA and Europe
- Study enrollment began in December 2021 and is ongoing in all study arms

Active sites

France

- Gustave Roussy, Villejuif Cedex
- Institut Bergonié, Bordeaux Cedex

Italy

- Istituto Europeo di Oncologia, Milan

Spain

- VHIO Vall d'Hebron Instituto de Oncología, Barcelona
- Hospital Universitario 12 de Octubre, Madrid
- South Texas Accelerated Research Therapeutics (START) Madrid - Hospital Fundacion Jimenez Diaz, Madrid
- Instituto Valenciano de Oncología, Valencia
- Institut Català d'Oncologia – Badalona, Barcelona

United States

- The University of Arizona Cancer Center, Tucson, Arizona
- Sarah Cannon Research Institute/ HealthONE Denver, Denver, Colorado
- BRCR Medical Center, Plantation, Florida
- Sarah Cannon Research Institute/Florida Cancer Specialists, Orlando, Florida
- The University of Chicago Medical Center, Chicago, Illinois
- Community Health Network, Indianapolis, Indiana
- Dana-Farber Cancer Institute, Boston, Massachusetts
- Massachusetts General Hospital, Boston, Massachusetts
- University of Michigan, Ann Arbor, Michigan

- Columbia University Herbert Irving Comprehensive Cancer Center, New York, New York
- Memorial Sloan Kettering Cancer Center, New York, New York
- Sarah Cannon Research Institute/ Tennessee Oncology, Nashville, Tennessee
- The University of Texas Southwestern Medical Center, Dallas, Texas
- The University of Texas MD Anderson Cancer Center, Houston, Texas
- Virginia Cancer Specialists Research Institute and NEXT Oncology, Virginia, Fairfax, Virginia
- The University of Wisconsin Carbone Cancer Center, Madison, Wisconsin

References

- Vasan N and Cantley LC. *Nat Rev Clin Oncol.* 2022;19:471–485.
- Millis SZ, et al. *JAMA Oncol.* 2016;2:1565–1573.
- Rugo HS, et al. *Ann Oncol.* 2020;31:1001–1010.
- Varkaris A, et al. *Cancer Discov.* 2024;14:240–257.
- Pazolli E, et al. *Mol Cancer Ther.* 2021;20 (12 Suppl.):P251.

Acknowledgments

The authors would like to thank the study participants and their families, and study investigators and research staff at the study sites. This study was sponsored by Relay Therapeutics, Inc. Medical writing support was provided by Bernadette Tynan, of BOLDSCIENCE Inc., funded by Relay Therapeutics.

Abbreviations

AE, adverse event; BC, breast cancer; BID, twice per day; BOIN, Bayesian Optimal Interval; CBR, clinical benefit rate; CDK4/6, cyclin-dependent kinases 4/6; CTCAE, Common Terminology Criteria for Adverse Events; ctDNA, circulating tumor DNA; DoR, duration of response; HbA1c, glycosylated hemoglobin; HER2–, human epidermal growth factor receptor 2-negative; HNSCC, head and neck squamous cell carcinoma; HR+, hormone receptor-positive; IC50, half-maximal inhibitory concentration; MTD, maximum tolerated dose; mTOR, mammalian target of rapamycin; mut, mutations; ORR, objective response rate; pAKT, phosphorylated AKT; PD, pharmacodynamics; PFS, progression-free survival; PI3K α , phosphatidylinositol 3-kinase alpha; *PIK3CA*, phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha; PK, pharmacokinetics; PO, by mouth; QD, once daily; RECIST v1.1, Response Evaluation Criteria in Solid Tumors version 1.1; RP2D, recommended Phase 2 dose; SEM, standard error of the mean; WT, wild type.

American Society of Clinical Oncology,
May 31–June 4, 2024, Chicago, IL, USA

Copies of this poster obtained through the Quick Response (QR) Code are for personal use only and may not be reproduced without permission from ASCO and the author of this poster. For more information, please contact: ClinicalTrials@relaytx.com.

