

Molecular profiling after short-term preoperative endocrine therapy to identify ER+/HER2+ early breast tumours with favourable prognosis

Milana A Bergamino Sirvén^{1,3}, Elena López-Knowles^{3,4}, Xixuan Zhu^{1,3}, Holly Tovey^{1,3}, Hui Xiao^{3,4}, Lucy S Kilburn¹, Anastasia Alataki^{3,4}, Chris Holcombe⁵, Anthony Skene⁶, Ian E. Smith^{3,4}, John F. Robertson⁷, Judith M Bliss^{1,3}, Eugene F Schuster^{3,4}, Mitch Dowsett^{3,4}, Maggie CU Cheang^{1,3}, on behalf of POETIC investigators. ¹Clinical Trials and Statistics Unit (ICR-CTSU)- London, UK, ²Institut Català d'Oncologia, Badalona, Spain. ³The Institute of Cancer Research, London, UK; ⁴Royal Marsden Hospital, London, UK ⁵Liverpool University Hospitals, Liverpool; ⁶Royal Bournemouth and Christchurch NHS Foundation Trust, Bournemouth, UK. ⁷The University of Nottingham, Nottingham, UK. *All authors contributed equally.

AIMS

We hypothesize that molecular changes (MoIC) in Estrogen Receptor positive (ER+)/ Human Epidermal Growth Factor Receptor Positive (HER2+) early Breast Cancer (BC) vary according to response to Aromatase Inhibitors (AI) and predict recurrence outcome.

Our aims are 1. to characterise MoIC under short endocrine therapy (ET) and 2. to evaluate their association with response to treatment and survival.


BACKGROUND

- Twenty per cent of all BC overexpresses HER2+ and 50% of them are also ER+.
- ER+/HER2+ BC is heterogeneous with different behavior and outcome compared to ER-/HER2+ BC.
- The impeded anti-proliferative response of ER+/HER2+ to ET has been considered negated by treatment with anti-HER2 therapies.
- Residual disease after anti-HER2 therapy remains at a higher risk of recurrence.
- Studying the MoIC of ER+/HER2+ BC in response to ET will provide clinical insights to treatment options.

MATERIALS & METHODS

POETIC is a phase III trial of post-menopausal women with ER+ early BC randomized 2:1 to 2-weeks of peri-operative AI (POAI) vs control, followed by standard-of-care.


Figure 1. Study design



Baseline and on-treatment RNA from formalin-fixed paraffin-embedded samples were gene expression profiled with BC360™ panel, including the intrinsic subtypes (IS) and 46 key biological pathway signatures in BC.


The association of molecular features with Ki67 and TTR were assessed.

Figure 2. Shifts of intrinsic subtype (IS) from baseline to on-treatment timepoints in both groups separately: A. POAI and B. Controls.



- IS shifting was significantly more prevalent in POAI (37%, 79/213) than in controls (14%, 14/100) (p < 0.001).
- In POAI most Luminal B (LumB) shifted to Luminal A (LumA; 79%, 59/75).

Figure 3. A. BC360 Biological signature expression changes; log2 fold change and changes in PAM50 IS correlation coefficient from baseline to on-treatment samples according to early biological response to AI by Ki67_{2wk}.



- There was an overall upregulation of tumor immunity-related signatures.
- Proliferation, Homologous Recombination Deficiency (HRD), TP53 mutational status and ER-signaling were downregulated (FDR<0.05).
- Gene expression changes were significantly higher in Low-Low and High-Low compared to High-High Ki67 categories.
- Controls showed downregulation of PDL1 and upregulation of hypoxia (FDR < 0.05).

RESULTS

Figure 4. Kaplan Meier curves for Time to Recurrence (TTR) in POAI according to on-treatment IS




Table 1. Multivariable cox regression model for Ki67_{2wk} adjusted by standard clinic-pathological variables

Variable	Factors	HR (95% CI)	P-value
On-treatment Ki67	-	1.03 (1.01 – 1.05)	0.01 *
Age	-	1.08 (1.02 – 1.14)	0.0054 **
Tumour size	-	1.52 (1.21 – 1.91)	0.0003 ***
Nodal status	N0 (ref) (n=111)	-	-
	N1-3 (n=62)	1.52 (0.39 – 3.59)	0.7563
	N4+ (n=36)	2.20 (0.84 – 5.77)	0.1097
Concordance Index = 0.765 (se = 0.057) Likelihood Ratio Test = 29.35 (p = 0.00002)			
AIC = 205.195			

- LumA_{2wk} was associated with better outcome compared to LumB_{2wk} as well as Ki67_{2wk} value as continuous variable.

Table 2. Multivariable cox regression model for the PAM50 intrinsic subtypes in POAI adjusted by standard clinic-pathological variables (age>70, post-surgery tumour size and nodal status) at A. Baseline and B. On-treatment.

A. Baseline IS in POAI			
Variable	Factors	HR (95% CI)	P-value
Baseline PAM50 IS	LumA (ref) (n=43)	-	-
	LumB (n=75)	1.21 (0.30 – 4.80)	0.7871
	HER2-E (n=89)	1.72 (0.47 – 6.24)	0.4091
	Basal (n=3)	4.84 (0.48 – 49.1)	0.1823
Age	-	1.07 (1.01 – 1.12)	0.0133 *
Tumour size	-	1.50 (1.20 – 1.88)	0.0004 ***
Nodal status	N0 (ref) (n=112)	-	-
	N1-3 (n=62)	1.14 (0.37 – 3.47)	0.8202
	N4+ (n=36)	2.41 (0.90 – 6.42)	0.0791
Concordance Index = 0.742 (se = 0.059) Likelihood Ratio Test = 25.61 (p = 0.0006)			
AIC = 212.944			

B. On-treatment IS in POAI			
Variable	Factors	HR (95% CI)	P-value
On-treatment PAM50 IS	LumA (ref) (n=110)	-	-
	LumB (n=14)	5.41 (1.50 – 19.5)	0.0099 **
	HER2-E (n=83)	2.41 (0.87 – 6.73)	0.0919
	Basal (n=3)	6.55 (0.76 – 56.8)	0.0882
Age	-	1.07 (1.02 – 1.13)	0.01125 *
Tumour size	-	1.53 (1.22 – 1.92)	0.0003 ***
Nodal status	N0 (ref) (n=112)	-	-
	N1-3 (n=62)	1.14 (0.38 – 3.46)	0.8179
	N4+ (n=36)	2.37 (0.89 – 6.34)	0.0858
Concordance Index = 0.789 (se = 0.053) Likelihood Ratio Test = 31.43 (p = 0.00005)			
AIC = 207.1192			

- On-treatment IS provides more prognostic information than baseline IS.

CONCLUSIONS

In ER+/HER2+ early BC:

- There is a clinical utility of peri-operative ET, guiding who would have good prognosis for adjuvant ET and who may need additional therapy.
- There is significant upregulation of immune and endocrine-related genes and downregulation of proliferation, HRD and TP53 mutational status genes in early AI responders.
- Most LumB BC shift to LumA. On-treatment IS provides more prognostic information than baseline IS.
- On-treatment Ki67_{2wk} value as continuous variable was associated with differential outcome in POAI.

ACKNOWLEDGEMENTS

We would like to thank all the POETIC participants and staff and also Nanostring for their assistance. POETIC is co-sponsored by The Royal Marsden NHS Foundation Trust and the ICR. POETIC is funded by CRUK (CRUK/07/015) and coordinated by the ICR-CTSU. We acknowledge NHS funding to the NIHR Biomedical Research Centre at The Royal Marsden and the ICR and also Breast Cancer Now. We acknowledge Fundación Martín Escudero and Rio Hortega contracts 2022/2023 CM22/00101 for Milana Bergamino's fellowship funding. Maggie C. U. Cheang reports co-inventor of PAM50 Bioclassifier, with royalties paid, advisory role for Veracyte and research funding support by Nanostring Technologies. MB declares advisory funding from Eisai, AstraZeneca, Pfizer, Novartis and travel expenses from Novartis and AstraZeneca.