



Germline NQO1c.559C>T as a biomarker of genetic susceptibility to Immune Checkpoints Blockade resistance

Jose Carlos Benitez^{1,2}, Javier Olivera³, Juan Luis Onieva^{4,5}, Manuel Cobo-Dols^{3,5}, Andres Mesas¹, Alberto Rios-Muñoz^{4,5}, Elisabeth Perez-Ruiz^{2,3}, Maria Garrido^{4,5}, Paula Vazquez¹, Beatriz Martinez^{2,3}, Alexandra Cantero^{2,3}, Andrea Gonzalez^{2,3}, Juan Zafra¹, Isabel Barragan^{1,5}, Antonio Rueda-Dominguez^{1,5}

¹Medical Oncology Department, Regional and Virgen de la Victoria University Hospitals, Malaga, Spain; ²Faculty of Medicine, University of Malaga, Malaga, Spain; ³Biomedical Research Institute of Malaga and BIONAND Nanomedicine Platform (BIMABIONAND Platform) Translational Research in Cancer Immunotherapy group, Malaga, Spain; ⁴Department of Radiation Oncology, Virgen de la Victoria University Hospital, Malaga; ⁵Group of Pharmacogenetics, Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden.

Background

- Tobacco is the main risk factor for developing lung cancer. Yet, some heavy smokers do not develop cancer while others do at young ages.
- The study of cancer immunology has led to the surge of immune checkpoint blockade (ICB), which transformed the treatment landscape of cancer pts.
- NQO1 has been associated with extreme phenotypes of high risk for the development of tobacco-induced NSCLC and with poor survival outcomes for xeroderphs.
- Development of biomarkers for ICB has been mainly focused on tumor-related factors while genetic susceptibility of the host has been poorly explored to date and some evidence suggests that genetic dysregulation may enable immune escape.

Methods

- From June 2016 to January 2023, 326 NSCLC pts from two main University Hospitals of Malaga (Spain) receiving ICB were prospectively enrolled, blood samples were gathered for all patients receiving ICB.
- We selected a representative set of 60 consecutive patients for an exome sequencing in order to characterize the germline contribution to resistance (progression within the first 6 months of treatment).
- DNA was extracted from patient's PBMC fraction and prepared for exome sequencing. The raw sequencing data was processed to remove low-quality reads and aligned to a reference genome using BWA. Base recalibration, germline variant calling and variant filtering was performed through GATK. The remaining variants CRAVAT (Cancer-Related Analysis of Variants Toolkit).
- Survival probabilities were estimated with the Kaplan-Meier method and survival curves were compared using the log-rank test. We explored the lung cancer related genes among our results.

Results

- We found 26 germline variants associated with overall survival (OS) and progression free survival (PFS), and 52 genes variants with response to ICB. Univariate analysis of the 8 most significant genes variants related to the three survival outcomes are shown in table 3. The variants were missense or involved a stop gain/loss.

josecarlos.benitez@ibima.eu
 @JCBenitezMD

*First author declare no conflict of interest.

Results

Table 1. Descriptive analysis of our 60 patients cohort selected for an exome sequencing in PBMC.

Variable	N	%	P
SEX			
Male	14	23.3	<.001
Female	46	76.7	
HISTOLOGICAL SUBTYPE			
Non-adenocarcinoma	16	26.7	<.001
Adenocarcinoma	44	73.3	
AGE AT DIAGNOSIS			
< 65	31	51.7	0.897
> 65	29	48.3	
STAGE AT DIAGNOSIS			
IA	1	1.7	<.001
IB	1	1.7	
IIA	2	3.3	<.001
IIB	2	3.3	
IIIA	4	6.7	<.001
IIIB	2	3.3	
IV	9	15.0	<.001
IVA	21	35.0	0.027
IVB	14	23.3	
IVC	3	5.0	<.001
ECOG			
0	10	16.7	<.001
1	45	75.0	
2	2	3.3	<.001
PDL1			
AN/D	3	5.0	<.001
[0,1]	21	35.0	0.092
[1,49]	20	33.3	
[50,100]	13	21.7	<.001
LPI			
AN/D	4	6.7	<.001
0	14	23.3	<.001
1	37	61.7	0.092
2	9	15.0	<.001

Note: %, is proportion ± 0.5
Median age at diagnosis was 64 (54-85) years, 76.6% were men, 44/60 (73.3%) were adenocarcinoma, PD-L1 was 21% in 55% of patients, 22% showed PD-L250%, 78% of patients reported advanced disease at diagnosis, 65% received ICB in combination with chemotherapy.




Figure 1. We found NQO1c.559C>T in 46.6% of pts (28/60). The variation was not associated with survival outcomes in the multivariate analysis (p>0.05). Kaplan-Meier curve of median PFS and OS for NQO1 c.559C>T genetic variant, comparing mutation (blue line) and wild type (red line) is shown in figure 1A and 1B respectively. However, we found a poor OS for non-ADC histology (Figure 1D) (19.7m vs 6m for mutated and WT, respectively; p<0.01). Multivariate analysis shown KRASL:565T>C, ZNF4c:488G>A and RREB1c:4315G>A as independent prognostic factors for OS, the two later for PFS as well. SLC17A3c:286G>A, SAMM5c:329A>G and RREB1c:4315G>A were independent predictive factors of response at 6 months in the Likelihood Ratio Test (LRT). Figure 2 shows the multivariate analysis for OS (figure 2A) and PFS (figure 2B) of main ICB prognostic factors

Figure 2

Variable	N	Hazard ratio	P
PDL1			
Negative	18	Reference	
Inter	17	0.47 (0.19, 1.17)	0.105
Positive	12	0.30 (0.11, 0.84)	0.022
Age_group			
<65	26	Reference	
>65	22	1.72 (0.76, 3.78)	0.177
LPI			
Negative	12	Reference	
Inter	27	1.04 (0.36, 3.01)	0.946
Positive	9	0.98 (0.27, 3.54)	0.980
Diagnosis_subtype			
Adenocarcinoma	35	Reference	
Non-Adeno	13	3.13 (1.38, 7.23)	0.006
T1_ECOC			
Adenocarcinoma	46	3.15 (1.15, 8.64)	0.026
NQO1			
Mut	20	Reference	
WT	28	0.91 (0.43, 1.94)	0.812

Variable	N	Hazard ratio	P
PDL1			
Negative	18	Reference	
Inter	17	0.36 (0.14, 0.94)	0.04
Positive	12	0.29 (0.11, 0.80)	0.02
Age_group			
<65	26	Reference	
>65	21	1.59 (0.76, 3.34)	0.22
LPI			
Negative	12	Reference	
Inter	26	0.79 (0.25, 2.45)	0.68
Positive	9	0.53 (0.22, 1.30)	0.78
Diagnosis_subtype			
Adenocarcinoma	34	Reference	
Non-Adeno	13	2.18 (0.90, 5.29)	0.09
T1_ECOC			
Adenocarcinoma	47	2.55 (1.06, 6.10)	0.04
NQO1			
Mut	20	Reference	
WT	27	1.11 (0.53, 2.33)	0.78

Gene	Variant	OS	PFS	LRT
SLC17A3	c.286G>A	0.0306	0.1128	0.0003
MYO10A	c.287G>T	0.0206	0.0700	0.0024
IRS1	c.1507C>G	0.0019	0.0146	0.0001
BAG1LNT4	c.280C>T	0.0810	0.0480	0.0001
SAMM5	c.329A>G	0.1088	0.0410	0.0001
ZNF4	c.488G>A	0.0607	0.0448	0.0001
RREB1	c.4315G>A	0.0148	0.0113	0.0001
ARAP1	c.346G>A	0.1660	0.1297	0.0001

Table 3. Multivariate analysis of the main gene variations in our sample. We can see in red color those mutations which seem to be statistically significant (p<0.05).

Conclusions

Exome sequencing analysis of germline variants could be a suitable methodology to identify suboptimal biomarkers of immune resistance in NSCLC patients. NQO1c.559C>T might help to identify poor prognosis non-ADC pts in the ICB setting.